

Product Data Sheet

Product Name: Regorafenib

Cat. No.: GC10111

Chemical Properties

Cas. No. 755037-03-7

Chemical Name 4-(4-(3-(4-chloro-3-(trifluoromethyl)phenyl)ureido)-3-fluorophenoxy)-N-methylpicolinamide

SMILES CNC(=O)C1=NC=CC(=C1)OC2=CC(=C(C=C2)NC(=O)NC3=CC(=C(C=C3)Cl)C(F)(F)F)C(F)(F)F


Formula $C_{21}H_{15}ClF_4N_4O_3$ M.Wt 482.82

Solubility $\geq 100\text{mg/mL}$ in DMSO Storage Desiccate at -20°C

General tips For obtaining a higher solubility, please warm the tube at 37°C and shake it in the ultrasonic bath for a while. Stock solution can be stored below -20°C for several months.

Shipping Condition Evaluation sample solution: ship with blue ice All other available size: ship with RT, or blue ice upon request.

Structure



Protocol

Cell experiment [1]:

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

Cell lines	NCI-H1703, NCI-H2170, SK-MES-1 cells
Preparation Method	Lung squamous cell carcinoma (LSCC) cell lines were exposed to different concentrations of Regorafenib for 24, 48, and 72h, 0-5 μ M for NCI-H1703 and 0-20 μ M for NCI-H2170 and SK-MES-1. Cell proliferation was measured by SRB assay.
Reaction Conditions	0-20 μ M; 24, 48, 72h
Applications	Regorafenib decreased the survival fractions of LSCC cells in a dose- and time-dependent manner.

Animal experiment [2]:

Animal models	Female athymic nu/nu mice
Preparation Method	Mice with orthotopic adrenal SK-N-SH neuroblastoma xenograft tumours were treated with vehicle or Regorafenib. Mice in group 1 were gavaged once daily with vehicle alone, while the other group of mice was fed once daily with Regorafenib at 30mg/kg. Both groups of mice were treated for 14 days and then euthanised by CO ₂ followed by cervical dislocation. Tumours were harvested, weighed and photographed.
Dosage form	30mg/kg; 14 days; p.o.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

Applications Regorafenib reduces xenograft tumour growth. The average final weight of the tumors in the control group was 2.16g, while the average final weight of the tumors in the Regorafenib-treated group was only 0.35g.

References:

- [1] Hu X, Wu L, Zhang Z, et al. The anti-tumor effect of regorafenib in lung squamous cell carcinoma in vitro[J]. Biochemical and Biophysical Research Communications, 2018, 503(2): 1123-1129.
- [2] Subramonian D, Phanhthilath N, Rinehardt H, et al. Regorafenib is effective against neuroblastoma in vitro and in vivo and inhibits the RAS/MAPK, PI3K/Akt/mTOR and Fos/Jun pathways[J]. British journal of cancer, 2020, 123(4): 568-579.

Background

Regorafenib is an orally active multi-kinase inhibitor with IC₅₀ values of 13/4.2/46, 22, 7, 1.5 and 2.5nM for VEGFR1/2/3, PDGFR β , Kit, RET and Raf-1, respectively^[1]. Regorafenib has good anti-tumor and anti-angiogenic activities^[2, 3].

In vitro, Regorafenib (0-20 μ M) treatment of lung squamous cell carcinoma (LSCC) cell lines (NCI-H1703, NCI-H2170, SK-MES-1 cells) for 24-72h, inhibited cell proliferation in a dose- and time-dependent manner and induced cell G0/G1 phase arrest^[4]. Regorafenib (2.5-10 μ M) treatment of HepG2 cells for 24-96h, inhibited cell growth in a dose- and time-dependent manner, induced cell apoptosis, and increased the level of phosphorylated c-Jun^[5].

In vivo, oral treatment of mice with adrenal SK-N-SH neuroblastoma orthotopically implanted with Regorafenib (30mg/kg) for 14 days, significantly inhibited tumor growth in the mice, with the average final weight of the tumor in the control group being 2.16g and the average final weight of the tumor in the Regorafenib-treated group being only 0.35g^[6]. Oral treatment of mice with metastatic colon cancer model with Regorafenib (30mg/kg) for 14 days, inhibited tumor growth and metastasis in the mice, significantly reduced tumor angiogenesis, and significantly reduced the number of infiltrating macrophages^[7].

References:

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

- [1] Wilhelm S M, Dumas J, Adnane L, et al. Regorafenib (BAY 73-4506): a new oral multikinase inhibitor of angiogenic, stromal and oncogenic receptor tyrosine kinases with potent preclinical antitumor activity[J]. International journal of cancer, 2011, 129(1): 245-255.
- [2] Chen C H, Hsu F T, Chen W L, et al. Induction of apoptosis, inhibition of MCL-1, and VEGF-A expression are associated with the anti-cancer efficacy of magnolol combined with regorafenib in hepatocellular carcinoma[J]. Cancers, 2021, 13(9): 2066.
- [3] Lim Y, Han S W, Yoon J H, et al. Clinical implication of anti-angiogenic effect of regorafenib in metastatic colorectal cancer[J]. PLoS One, 2015, 10(12): e0145004.
- [4] Hu X, Wu L, Zhang Z, et al. The anti-tumor effect of regorafenib in lung squamous cell carcinoma in vitro[J]. Biochemical and Biophysical Research Communications, 2018, 503(2): 1123-1129.
- [5] Carr B I, Cavallini A, Lippolis C, et al. Fluoro-Sorafenib (Regorafenib) effects on hepatoma cells: growth inhibition, quiescence, and recovery[J]. Journal of cellular physiology, 2013, 228(2): 292-297.
- [6] Subramonian D, Phanhthilath N, Rinehardt H, et al. Regorafenib is effective against neuroblastoma in vitro and in vivo and inhibits the RAS/MAPK, PI3K/Akt/mTOR and Fos/Jun pathways[J]. British journal of cancer, 2020, 123(4): 568-579.
- [7] Abou-Elkacem L, Arns S, Brix G, et al. Regorafenib inhibits growth, angiogenesis, and metastasis in a highly aggressive, orthotopic colon cancer model[J]. Molecular cancer therapeutics, 2013, 12(7): 1322-1331.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA