

Product Data Sheet

Product Name: ODQ
Cat. No.: GC10279

Chemical Properties

Cas. No. 41443-28-1

Chemical Name 1H-[1,2,4]oxadiazolo[4,3-a]quinoxalin-1-one

SMILES O=C1ON=C2N1C3=CC=CC=C3N=C2


Formula $C_9H_5N_3O_2$ M.Wt 187.16

Solubility DMF: 30 mg/ml, DMF:PBS (pH 7.2) (1:1): .1 mg/ml, DMSO: 15 mg/ml, Ethanol: .5 mg/ml
Storage Store at -20°C

General tips For obtaining a higher solubility, please warm the tube at 37 °C and shake it in the ultrasonic bath for a while. Stock solution can be stored below -20°C for several months.

Shipping Condition Evaluation sample solution: ship with blue ice. All other available size: ship with RT, or blue ice upon request.

Structure



Protocol

Cell experiment [1]:

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

Cell lines NBL-6 cells

Preparation Method Cells were infected with EqHV-8 at an MOI of 0.1 for 1 h, followed by the treatment of cells with or without CoPP (100 μ M) in the presence or absence of ODQ (10 μ M).

Reaction
Conditions 10 μ M

Applications ODQ partially reversed the anti-viral effect of cobalt-protoporphyrin (CoPP) by EqHV-8 progeny virus generation.

Animal experiment [2]:

Animal models C57BL/6J mice(LLC tumor model)

Preparation Method Mice received oral administration of 2.5mg/kg Fruquintinib and intraperitoneal injection of 6mg/kg ODQ, both starting from day 6 post-tumor injection, and tumor analysis at day 14 post-tumor injection.

Dosage form i.p; 6mg/kg; 8 days

Applications ODQ treatment alone reduced tumor growth, the combination of ODQ and Fruquintinib further decreased tumor size.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

References:

- [1]. Wang T, Li S, et.al. Heme oxygenase-1 is an equid alphaherpesvirus 8 replication restriction host protein and suppresses viral replication via the PKC α /ERK1/ERK2 and NO/cGMP/PKG pathway. Microbiol Spectr. 2024 Apr 2;12(4):e0322023. doi: 10.1128/spectrum.03220-23. Epub 2024 Mar 5. PMID: 38441979; PMCID: PMC10986571.
- [2]. Zhu J, Yang W, et.al. Pericyte signaling via soluble guanylate cyclase shapes the vascular niche and microenvironment of tumors. EMBO J. 2024 Apr;43(8):1519-1544. doi: 10.1038/s44318-024-00078-5. Epub 2024 Mar 25. PMID: 38528180; PMCID: PMC11021551.

Background

ODQ is a soluble guanylyl cyclase (sGC) inhibitor that binds sGC in a competitive but irreversible manner [1-2].

ODQ(10 μ M) partially reversed the antiviral effect of CoPP by inhibiting Soluble guanylyl cyclase (sGC) [3]. ODQ(20 μ M) reduces the ability of Naringenin (NAR) to promote the osteogenic differentiation of human periodontal ligament stem cells (hPDLSCs) [4]. ODQ(30 μ M) significantly induced apoptosis of NCI-H2452 cells [5].

ODQ(i.p; 6mg/kg; 8 days) treatment alone can slow the growth rate of tumors, the combination of ODQ and Fruquintinib further decreased tumor size[6]. ODQ(25-100 μ g/mL) can eliminate the protective effect of Zhenwu Decoction (ZWD) on myocardial hypertrophy by inhibiting sGC-cGMP-PKG signaling pathway in zebrafish [7].

References:

- [1]. Feelisch M, Kotsonis P, et.al. The soluble guanylyl cyclase inhibitor 1H-[1,2,4]oxadiazolo[4,3,-a]quinoxalin-1-one is a nonselective heme protein inhibitor of nitric oxide synthase and other cytochrome P-450 enzymes involved in nitric oxide donor bioactivation. Mol Pharmacol. 1999 Aug;56(2):243-53. doi: 10.1124/mol.56.2.243. PMID: 10419542.
- [2]. Schrammel A, Behrends S, et.al. Characterization of 1H-[1,2,4]oxadiazolo[4,3-a]quinoxalin-1-one as a heme-site inhibitor of nitric oxide-sensitive guanylyl cyclase. Mol Pharmacol. 1996 Jul;50(1):1-5. PMID: 8700100.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

- [3]. Wang T, Li S, et.al. Heme oxygenase-1 is an equid alphaherpesvirus 8 replication restriction host protein and suppresses viral replication via the PKC β /ERK1/ERK2 and NO/cGMP/PKG pathway. Microbiol Spectr. 2024 Apr 2;12(4):e0322023. doi: 10.1128/spectrum.03220-23. Epub 2024 Mar 5. PMID: 38441979; PMCID: PMC10986571.
- [4]. Li S, Xiong Z, et.al. Naringenin modulates the NO-cGMP-PKG signaling pathway by binding to AKT to enhance osteogenic differentiation in hPDLSCs. Int J Mol Med. 2024 Aug;54(2):67. doi: 10.3892/ijmm.2024.5391. Epub 2024 Jun 28. PMID: 38940332; PMCID: PMC11232664.
- [5]. Ronald Fiscus, et al. ODQ, an inhibitor of soluble guanylyl cyclase (nitric oxide-activated enzyme), enhances the pro-apoptotic effects of cisplatin in human mesothelioma cells. Cancer Res May 1 2007 (67) (9 Supplement) LB-43.
- [6]. Zhu J, Yang W, et.al. Pericyte signaling via soluble guanylate cyclase shapes the vascular niche and microenvironment of tumors. EMBO J. 2024 Apr;43(8):1519-1544. doi: 10.1038/s44318-024-00078-5. Epub 2024 Mar 25. PMID: 38528180; PMCID: PMC11021551.
- [7]. Chen L, Zhou X, et.al. Zhenwu decoction ameliorates cardiac hypertrophy through activating sGC (soluble guanylate cyclase) - cGMP (cyclic guanosine monophosphate) - PKG (protein kinase G) pathway. J Ethnopharmacol. 2023 Jan 10;300:115705. doi: 10.1016/j.jep.2022.115705. Epub 2022 Sep 12. PMID: 36099983.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA