

Product Data Sheet

Product Name: OSU-2S TFA

Cat. No.: GC81880

Chemical Properties

Cas. No.

Formula $C_{23}H_{38}F_3NO_4$ M.Wt 449.55

Solubility DMSO: 100 mg/mL (222.44 mM; Need ultrasonic) Storage Store at 4°C, sealed storage, away from moisture and light

General tips For obtaining a higher solubility, please warm the tube at 37 °C and shake it in the ultrasonic bath for a while. Stock solution can be stored below -20°C for several months.

Shipping Condition Evaluation sample solution : ship with blue ice All other available size: ship with RT, or blue ice upon request.

Structure



Background

OSU-2S TFA is a non-immunosuppressive derivative of Fingolimod hydrochloride (FTY720). It is an orally active, blood-brain barrier-permeable allosteric activator of PP2A that also activates PKC δ ; it restores PP2A activity by displacing the inhibitory protein SET, thereby inducing ROS - PKC δ - caspase-3 -dependent apoptosis, c-Myc degradation, cell cycle arrest, and differentiation. OSU-2S TFA is used in research on diseases such as leukemia, liver cancer, lymphoma, and lung cancer [1] [2] [3] [4] [5] [6] [7].

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

In Vivo, OSU-2S (2.5–10 mg/kg; intraperitoneal injection; every other day for 35 days) TFA inhibits tumor growth in an A549 xenograft model, is well tolerated, and downregulates Ki-67 and VEGF^[1].

In Vitro, OSU-2S (2-6 μ M; 24 h) TFA exhibits concentration-dependent cytotoxicity against canine B-cell lymphoma cell lines (CLBL-1 and 17-71)^[7]. The cytotoxicity of OSU-2S (3 μ M; 24 h) TFA in canine B-cell lymphoma cell lines is partially mediated by ROS^[7]. OSU-2S (2-4 μ M; 24 h) TFA induces ROS production and mediates cytotoxicity in primary cells from canine spontaneous B-cell lymphoma^[7]. OSU-2S (24 h) TFA downregulates PGC1 α and inhibits mitochondrial biogenesis via PP2A-dependent FOXO1 dephosphorylation^[6]. OSU-2S TFA reduces mitochondrial mass and ATP production, and inhibits the expression of genes associated with mitochondrial biogenesis^[6]. OSU-2S (2 μ M; 24 h) TFA induces apoptosis in primary CLL cells^[4]. OSU-2S TFA activates PP2A in CLL cells^[4]. OSU-2S (5 h) TFA induces SHP1 S591 phosphorylation via a PKC-dependent mechanism^[4]. OSU-2S (5-24 h) TFA -induced pSHP1 S591 levels correlate with CLL cell death ^[4]. The cytotoxicity of OSU-2S (2 μ M; 24 h) TFA is partially dependent on PKC^[4]. OSU-2S (8 μ M; 4 h) TFA activates PKC in CLL cells^[4]. OSU-2S (8 μ M; 16 h) TFA extensively regulates gene expression in CLL cells, affecting processes such as cell growth, proliferation, death, and the cell cycle^[4]. OSU-2S (8 μ M; 16-24 h) TFA downregulates TCL1A expression in CLL cells^[4]. OSU-2S (5 μ M; 24 h) TFA selectively kills CLL cells without affecting normal B cells via ROR1-targeted delivery^[4]. OSU-2S (2-8 μ M; 24 h) TFA exhibits potent cytotoxicity against MCL cell lines^[3]. OSU-2S (2-8 μ M; 24 h) TFA induces apoptosis in MCL cell lines^[3]. OSU-2S (4-5 μ M; 24 h) TFA induces CD74 expression on the surface of MCL cells^[3]. OSU-2S (4 μ M; 24 h) TFA enhances the cytotoxicity of Milatuzumab against MCL cells by upregulating CD74^[3]. OSU-2S (2-10 μ M; 48 h) TFA demonstrates anti-proliferative activity against various HCC cell lines^[2]. OSU-2S (1.25-2.5 μ M; 48 h) TFA enhances the anti-proliferative effect of Sorafenib on HCC cells, exhibiting a synergistic effect^[2]. OSU-2S (1.25-2.5 μ M; 48 h) TFA enhances Sorafenib-induced apoptosis in HCC cells^[2]. OSU-2S (1.25-2.5 μ M; 48 h) TFA enhances the anti-HCC effect of Sorafenib via PKC δ and ERK pathways^[2]. OSU-2S (1.25-2.5 μ M; 8-24 h) TFA in combination with Sorafenib inhibits HCC cell migration and invasion^[2]. The activity of OSU-2S (1.25-7.5 μ M) TFA against HCC cells is influenced by p53 status; p53 restoration enhances

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

sensitivity^[2]. OSU-2S (1-10 μ M; 24 h) TFA inhibits A549 proliferation, with an IC₅₀ of 4.432 μ M^[1]. OSU-2S (1.5-6 μ M; 14 days) TFA inhibits colony formation in A549 cells^[1]. OSU-2S (1.5-6 μ M; 12-24 h) TFA inhibits A549 cell migration^[1]. OSU-2S (1.5-6 μ M; 24 h) TFA downregulates p-AURKA and S1PR1 protein expression in A549 cells^[1].

References:

- [1]. Han M, et al. Evaluation of the Efficacy of OSU-2S in the Treatment of Non-Small-Cell Lung Cancer and Screening of Potential Targets of Action. *Pharmaceuticals (Basel)*. 2024 May 1;17(5):582.
- [2]. Omar HA, et al. OSU-2S/Sorafenib Synergistic Antitumor Combination against Hepatocellular Carcinoma: The Role of PKC δ /p53. *Front Pharmacol*. 2016 Nov 30;7:463.
- [3]. Mani R, et al. ROR1-targeted delivery of OSU-2S, a nonimmunosuppressive FTY720 derivative, exerts potent cytotoxicity in mantle-cell lymphoma in vitro and in vivo. *Exp Hematol*. 2015 Sep;43(9):770-4.e2.
- [4]. Mani R, et al. Tumor antigen ROR1 targeted drug delivery mediated selective leukemic but not normal B-cell cytotoxicity in chronic lymphocytic leukemia. *Leukemia*. 2015 Feb;29(2):346-55.
- [5]. Xie Z, et al. Pharmacokinetics and Tolerability of the Novel Non-immunosuppressive Fingolimod Derivative, OSU-2S, in Dogs and Comparisons with Data in Mice and Rats. *AAPS J*. 2020 Jul 16;22(4):92.
- [6]. Khalil SM, et al. Protein Phosphatase 2A Activation Dependent Down Regulation of PGC1-Alpha and FOXO1 Phosphorylation with OSU-2S in Human Lymphoma Cells. *Blood*. 2019;134(Supplement_1):3977.
- [7]. Mani R, et al. Non-immunosuppressive FTY720-derivative OSU-2S mediates reactive oxygen species-mediated cytotoxicity in canine B-cell lymphoma. *Vet Comp Oncol*. 2017 Sep;15(3):1115-1118.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA