

Product Name: NRX-0492
Cat. No.: GC73434

Cas. No. 2416130-57-7


Formula	C ₄₃ H ₅₁ N ₁₁ O ₆	M.Wt	817.94
---------	--	------	--------

Solubility	DMSO : 200 mg/mL (244.52 mM; Need ultrasonic)	Storage	-20°C
------------	---	---------	-------

General tips	For obtaining a higher solubility , please warm the tube at 37 °C and shake it in the ultrasonic bath for a while. Stock solution can be stored below -20°C for several months.
--------------	---

Shipping Condition	Evaluation sample solution : ship with blue ice All other available size: ship with RT , or blue ice upon request.
--------------------	--

Structure



NRX-0492 is an orally active and potent degrader of BTK. NRX-0492 catalyzes ubiquitylation and proteasomal degradation of BTK with $DC_{50} \leq 0.2$ nM and $DC_{90} \leq 0.5$ nM, respectively. NRX-0492 inhibits B-cell receptor (BCR)-mediated signaling, transcriptional programs, and chemokine secretion.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

Moreover, NRX-0492 also links a noncovalent BTK-binding domain to Cereblon. Cereblon is an adaptor protein of the E3 ubiquitin ligase complex.

References:

[1]. Zhang D, et al. NRX-0492 degrades wild-type and C481 mutant BTK and demonstrates in vivo activity in CLL patient-derived xenografts. *Blood*. 2023 Mar 30;141(13):1584-1596.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA