

Product Data Sheet

Product Name: NC03
Cat. No.: GC27959

Chemical Properties

Cas. No. 568558-66-7

Chemical Name Acetamide, N-(2,3-dihydro-1,4-benzodioxin-6-yl)-2-[[5-(3,4,5-trimethoxyphenyl)-1,3,4-oxadiazol-2-yl]thio]-

SMILES O=C(CSC1=NN=C(C2=CC(OC)=C(OC)C(OC)=C2)O1)NC3=CC=C4OCCOC4=C3


Formula $C_{21}H_{21}N_3O_7S$ M.Wt 459.47

Solubility DMSO : 100 mg/mL (217.64 mM; Need ultrasonic) Storage -20°C

General tips For obtaining a higher solubility , please warm the tube at 37 °C and shake it in the ultrasonic bath for a while. Stock solution can be stored below -20°C for several months.

Shipping Condition Evaluation sample solution : ship with blue ice All other available size: ship with RT , or blue ice upon request.

Structure



Background

NC03 is a PI4K2A inhibitor and reducing agent. NC03 inhibits the activity of PI4K2A, thereby reducing the production of PI4P in the cellular compartments where PI4K2A functions. NC03 rapidly induces a decrease in PI4P pools in the Golgi and endosomal compartments. NC03 reduces PI4P

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

production in Rab7-positive late endosomes^{[1][2][3]}.

In Vitro, NC03 (15-76 μ M; 1 hour) potently inhibits the activity of bacterially expressed human PI4K2A enzyme in COS-7 cells^[1].

NC03 (10 μ M; 10 min pre-incubation, 1.5 h kinase reaction) inhibits the activity of palmitoylated PI4K2A enzyme isolated from COS-7 cells when used at 10 μ M^[1].

NC03 (1-10 μ M; 20 min pre-incubation, 30 min permeabilized cell incubation) inhibits PI4K2A-mediated PI4P synthesis in permeabilized COS-7 cells, with secondary effects on PI(4,5)P₂ levels^[1].

NC03 (10 μ M; up to 10 min) rapidly reduces Golgi and endosomal PI4P pools in live COS-7 cells when used at 10 μ M^[1].

NC03 (10-30 μ M; 30 min) decreases Rab7-positive late endosomal PI4P levels in intact HEK-AT1 cells in a concentration-dependent manner, with 30 μ M showing greater efficacy than 10 μ M^[1].

References:

- [1]. Kumari S, et al. Enhancement of aqueous solubility and oral bioavailability of nelfinavir by complexation with β -cyclodextrin[J]. Tropical Journal of Pharmaceutical Research, 2015, 14(8): 1333-1340.
- [2]. Hofacker A F, et al. Mercury mobilization in a flooded soil by incorporation into metallic copper and metal sulfide nanoparticles[J]. Environmental science & technology, 2013, 47(14): 7739-7746.
- [3]. Sengupta N, et al. A large scale high-throughput screen identifies chemical inhibitors of phosphatidylinositol 4-kinase type II alpha. J Lipid Res. 2019;60(3):683-693.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA