

Product Data Sheet

Product Name: Edoxudine (EUDR)

Cat. No.: GC32148

Chemical Properties

Cas. No. 15176-29-1

SMILES OC[C@@H]1[C@H](C[C@H](N2C(NC(C(CC)=C2)=O)=O)O1)O


Formula $C_{11}H_{16}N_2O_5$ M.Wt 256.26

Solubility DMSO : ≥ 125 mg/mL (487.79 mM) Storage Store at -20°C

General tips For obtaining a higher solubility , please warm the tube at 37°C and shake it in the ultrasonic bath for a while. Stock solution can be stored below -20°C for several months.

Shipping Condition Evaluation sample solution : ship with blue ice All other available size: ship with RT , or blue ice upon request.

Structure



Protocol

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

Cell experiment [1]:

Cell lines Human foreskin fibroblast (HFF)

Preparation Method 24h prior to assay, human foreskin fibroblasts (HFFs) were seeded in 6-well plates at a concentration of 2.5×10^4 cells per well in MEM containing 10% FBS. The media from the wells was then aspirated with different concentrations of Edoxudine (1, 10, 100, and 1000 μ M). The cells were incubated in a CO₂ incubator at 37°C for 72h. The media-drug solution was removed, and the cells were washed. 1ml of 0.25% trypsin was added to each well and incubated until the cells started to come off of the plate. The cell-media mixture was then pipetted up and down vigorously to break up the cell suspension, followed by cell counting.

Reaction Conditions 1, 10, 100, and 1000 μ M; 72h

Applications Edoxudine treatment decreased the cell viability of HFFs in a dose-dependent manner.

References:

[1] Kumar R, Wiebe L I, Knaus E E. Synthesis and antiviral activity of novel 5-(1-azido-2-haloethyl) and 5-(1-azido-, amino-, or methoxyethyl) analogs of 2'-deoxyuridine[J]. Journal of medicinal chemistry, 1993, 36(17): 2470-2474.

Background

Edoxudine (EUDR) is a deoxy-thymidine analog shown to be effective against herpes simplex virus type 1 and type 2^[1]. Edoxudine has been shown in vertebrate systems to be phosphorylated preferentially by the thymidine kinase and incorporated into viral DNA, and inhibiting viral DNA synthesis^[2]. Edoxudine disrupts the outer wall of *Klebsiella pneumoniae* and damages the protective effect of the LPS layer, making the cell membrane of *Klebsiella pneumoniae* accessible to

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

antibacterial effectors^[3]. Edoxudine has been widely used as a lead compound for the development of drugs targeting the surface protein of monkeypox virus E8 ^[4].

In vitro, Edoxudine treatment for 72 hours significantly inhibited the proliferation of human foreskin fibroblasts (HFFs), with an IC₅₀ value of 115µM^[5].

References:

- [1] Sharma A, Tiwari V, Sowdhamini R. Computational search for potential COVID-19 drugs from FDA-approved drugs and small molecules of natural origin identifies several anti-virals and plant products[J]. Journal of biosciences, 2020, 45(1): 100.
- [2] Mazzacano C A, Fallon A M. Evaluation of a viral thymidine kinase gene for suicide selection in transfected mosquito cells[J]. Insect molecular biology, 1995, 4(2): 125-134.
- [3] Ifrid E, Ouertatani-Sakouhi H, Jauslin T, et al. 5-ethyl-2'-deoxyuridine fragilizes Klebsiella pneumoniae outer wall and facilitates intracellular killing by phagocytic cells[J]. Plos one, 2022, 17(10): e0269093.
- [4] Podduturi S, Vemula D, Singothu S, et al. In-silico investigation of E8 surface protein of the monkeypox virus to identify potential therapeutic agents[J]. Journal of Biomolecular Structure and Dynamics, 2024, 42(16): 8242-8255.
- [5] Kumar R, Wiebe L I, Knaus E E. Synthesis and antiviral activity of novel 5-(1-azido-2-haloethyl) and 5-(1-azido-, amino-, or methoxyethyl) analogs of 2'-deoxyuridine[J]. Journal of medicinal chemistry, 1993, 36(17): 2470-2474.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA