

## Product Data Sheet

Product Name: Myristic Acid-d<sub>2</sub>

Cat. No.: GC47713

### Chemical Properties

Cas. No. 30719-21-2


SMILES OC(C([2H])([2H])CCCCCCCCCCCC)=O

Formula C<sub>14</sub>H<sub>26</sub>D<sub>2</sub>O<sub>2</sub> M.Wt 230.4

Solubility DMF: 15 mg/ml, DMSO: 10 mg/ml, Ethanol: 15 mg/ml  
 Storage Store at -20°C

General For obtaining a higher solubility, please warm the tube at 37 °C and shake it in the tips ultrasonic bath for a while. Stock solution can be stored below -20°C for several months.

Shipping Evaluation sample solution : ship with blue ice All other available size: ship with RT, or Condition blue ice upon request.



### Background

Myristic acid-d<sub>2</sub> is intended for use as an internal standard for the quantification of myristic acid by GC- or LC-MS. Myristic acid is a 14-carbon saturated fatty acid. It is incorporated into myristoyl coenzyme A (myristoyl-CoA) and transferred by N-myristoyltransferase to the N-terminal glycine of certain proteins either during translation to modify protein activity or post-translationally in apoptotic cells.<sup>1,2</sup>

1. Bhatnagar, R.S., Fütterer, K., Waksman, G., et al. The structure of myristoyl-CoA: Protein N-myristoyltransferase. *Biochim. Biophys. Acta* 1441(2-3):162-172 (1999)  
 2. Martin, D.D.O., Beauchamp, E., and Berthiaume, L.G. Post-translational myristoylation: Fat matters in cellular life and death. *Biochimie* 93(1):18-31 (2011)

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: [tech@glpbio.com](mailto:tech@glpbio.com)

Address: 10292 Central Ave. #205, Montclair, CA, USA