

Product Data Sheet

Product Name: Zilpaterol

Cat. No.: GC13333

Chemical Properties

Cas. No. 119520-05-7

Chemical Name (6R,7R)-*rel*-4,5,6,7-tetrahydro-7-hydroxy-6-[(1-methylethyl)amino]-imidazo[4,5,1-jk][1]benzazepin-2(1H)-one

SMILES O=C1NC2=CC=CC3=C2N1CC[C@H](NC(C)C)[C@H]3O


Formula $C_{14}H_{19}N_3O_2$ M.Wt 261.3

Solubility ≤ 1 mg/ml in DMSO; 0.1mg/ml in dimethyl formamide Storage Store at -20°C

General tips For obtaining a higher solubility, please warm the tube at 37°C and shake it in the ultrasonic bath for a while. Stock solution can be stored below -20°C for several months.

Shipping Condition Evaluation sample solution: ship with blue ice All other available size: ship with RT, or blue ice upon request.

Structure



Background

Zilpaterol is a β -adrenergic receptor agonist.

β -adrenergic receptors, prototypic G-protein-coupled receptors, play a key role in regulating numerous physiological processes.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

In vitro: Previous study found that there was no effect of zilpaterol (ZH) on thymidine incorporation into DNA. ZH at 1 μ M could decrease β 1-AR mRNA, and 0.01 and 1 μ M ZH decreased β 2-AR and β 3-AR mRNA in myoblasts. There was no effect of ZH on the β -AR or IGF-I gene expression in fused myotube cultures. The β 2-AR antagonist ICI-118, 551 attenuated the effect of ZH to reduce expression of β 1- and β 2-AR mRNA. There was no effect of ICI-118, 551 or ZH on β 3-AR or IGF-I. In addition, the total lipid synthesis from acetate was increased by ZH in bovine subcutaneous adipose tissue explants in the absence of theophylline [1].

In vivo: FDA has approved label claims of increased rate of BW gain, improved feed efficiency, and increased carcass leanness. The feeding of ZH for 20 to 40 d with a 3-d withdrawal resulted in significantly increased ADG. Such increases equate to an average of 9 kg heavier BW in ZH-fed steers. Moreover, hot carcass weight has been shown to increase to a larger degree compared with BW, with an average improvement of 15 kg [2].

Clinical trial: So far, no clinical study has been conducted.

References:

- [1] Miller, E. K., Chung, K.Y., Hutcheson, J.P., et al. Zilpaterol hydrochloride alters abundance of β -adrenergic receptors in bovine muscle cells but has little effect on de novo fatty acid biosynthesis in bovine subcutaneous adipose tissue explants. *Journal of Animal Science* 90(4), 1317-1327 (2012).
- [2] Delmore, R. J., Hodgen, J.M. and Johnson, B.J. Perspectives on the application of zilpaterol hydrochloride in the United States beef industry. *Journal of Animal Science* 88(8), 2825-2828 (2010).

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA