

Product Data Sheet

Product Name: Ketoisovaleric acid

Cat. No.: GC13084

Chemical Properties

Cas. No. 759-05-7

Chemical Name 3-methyl-2-oxobutanoic acid

SMILES CC(C(C(=O)=O)=O)C


Formula $C_5H_8O_3$ M.Wt 116.12

Solubility $\geq 11.6\text{mg/mL}$ in DMSO Storage Store at -20°C

General tips For obtaining a higher solubility, please warm the tube at 37°C and shake it in the ultrasonic bath for a while. Stock solution can be stored below -20°C for several months.

Shipping Condition Evaluation sample solution: ship with blue ice. All other available size: ship with RT, or blue ice upon request.

Structure



Protocol

Cell experiment [1]:

Cell lines Human primary lymphocyte

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

Peripheral venous blood was obtained from healthy adult individuals, and the cell was separated by the Ficoll-Hypaque (Pharmacia) gradient method.

Preparation Method Lymphocytes (1×10^5 cells) were incubated at 37°C, a humidified atmosphere of 5% CO₂ and 95% air in the presence of phytohaemagglutinin and 20% autologous serum. The cells were treated with 5mM Ketoisovaleric acid within 24h.

Reaction
Conditions 5mM; 24h

Applications Ketoisovaleric acid moderately suppressed lymphocyte DNA synthesis and caused an immunosuppression in vitro.

Animal experiment [2]:

Animal models Male Wistar rats

Preparation Method 10min before training, a 30-gauge needle was inserted into the guide cannula, protruding 1mm from the tip of the cannula in order to administer a bilateral infusion of 2µl Ketoisovaleric acid into the dorsal hippocampus was administered to male rat, which affected rat performance in the inhibitory avoidance task, suggesting disruption of acquisition.

Dosage form 5µmol/2µl; 10min; intrahippocampal injection

Applications Ketoisovaleric acid affected rat performance in the inhibitory avoidance task.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

References:

- [1] SCHLOTTFELDT J, BLAZINA L R, WANNMACHER C M, et al. The effect of organic acids on phytohaemagglutinin-activated proliferation of human lymphocytes in vitro [J]. Int J Immunopharmacol, 1995, 17(3): 175-82.
- [2] DE CASTRO VASQUES V, DE BOER M A, DILIGENTI F, et al. Intrahippocampal administration of the alpha-keto acids accumulating in maple syrup urine disease provokes learning deficits in rats [J]. Pharmacol Biochem Behav, 2004, 77(1): 183-90.

Background

Ketoisovaleric acid (KIV) is as a precursor in leucine and valine synthesis^[1]. Ketoisovaleric acid can serve as an initial compound in vitamin B5 biosynthesis of Escherichia coli^[2]. Long-term high levels of Ketoisovaleric acid are associated with maple syrup urine disease (MSUD)^[3, 4].

Ketoisovaleric acid (1 to 5mM; 1h) significantly reduced the content of nonenzymatic antioxidants in the cerebral cortex in vitro, thereby decreasing the antioxidant responsiveness of cortical cells^[3]. Ketoisovaleric acid (5mM; 24h) moderately suppressed lymphocyte DNA synthesis, which caused an in vitro immunosuppression^[5].

Ketoisovaleric acid (5μmol/2μl; intrahippocampal injection; 10min) was administered to male rat, which provoked rat learning deficits in the inhibitory avoidance task^[6]. Ketoisovaleric acid (8μmol/2μl; intrastriatal administration; 20min) could induce seizures in rats^[7].

References:

- [1] NSHIMIYIMANA P, LIU L, DU G. Engineering of L-amino acid deaminases for the production of alpha-keto acids from L-amino acids [J]. Bioengineered, 2019, 10(1): 43-51.
- [2] CHASSAGNOLE C, DIANO A, LETISSE F, et al. Metabolic network analysis during fed-batch cultivation of Corynebacterium glutamicum for pantothenic acid production: first quantitative data and analysis of by-product formation [J]. J Biotechnol, 2003, 104(1-3): 261-72.
- [3] BRIDI R, BRAUN C A, ZORZI G K, et al. alpha-keto acids accumulating in maple syrup urine disease stimulate lipid peroxidation and reduce antioxidant defences in cerebral cortex from young rats [J]. Metab Brain Dis, 2005, 20(2): 155-67.
- [4] RABELO F, LEMOS I D S, DAL TOE C P, et al. Acute effects of intracerebroventricular

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

administration of alpha-ketoisocaproic acid in young rats on inflammatory parameters [J]. Metab Brain Dis, 2023, 38(5): 1573-9.

[5] SCHLOTTFELDT J, BLAZINA L R, WANNMACHER C M, et al. The effect of organic acids on phytohaemagglutinin-activated proliferation of human lymphocytes in vitro [J]. Int J Immunopharmacol, 1995, 17(3): 175-82.

[6] DE CASTRO VASQUES V, DE BOER M A, DILIGENTI F, et al. Intrahippocampal administration of the alpha-keto acids accumulating in maple syrup urine disease provokes learning deficits in rats [J]. Pharmacol Biochem Behav, 2004, 77(1): 183-90.

[7] COITINHO A S, DE MELLO C F, LIMA T T, et al. Pharmacological evidence that alpha-ketoisovaleric acid induces convulsions through GABAergic and glutamatergic mechanisms in rats [J]. Brain research, 2001, 894(1): 68-73.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA