

Product Data Sheet

Product Name: 18 α -Glycyrrhetic acid

Cat. No.: GC35061

Chemical Properties

Cas. No. 1449-05-4

SMILES CC1(C)[C@@H](O)CC[C@@]2(C)[C@@]1([H])CC[C@]([C@@](CC[C@]3(C)[C@]4([H])C[C@@](C)(C(O)=O)CC3)(C)C4=C5)(C)[C@]2([H])C5=O

Formula C₃₀H₄₆O₄

M.Wt 470.68


Solubility DMF: 13 mg/ml, DMSO: 16 mg/ml, Ethanol: 20 mg/ml, Ethanol: PBS (pH 7.2) (1:7): 0.13 mg/ml

Storage Store at -20°C

General tips For obtaining a higher solubility, please warm the tube at 37 °C and shake it in the ultrasonic bath for a while. Stock solution can be stored below -20°C for several months.

Shipping Condition Evaluation sample solution: ship with blue ice. All other available size: ship with RT, or blue ice upon request.

Structure



Protocol

Cell experiment [1]:

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

Cell lines	Human HSC cell line
Preparation Method	2×10^4 cells per well were seeded onto 24-well plates and incubated overnight. After starvation for 24h, the cells were subsequently stimulated with 10% FBS with or without the presence of 18 α -Glycyrrhetic acid. The number of cells was determined by computer-equipped cell counter 72h after treated with 18 α -Glycyrrhetic acid.
Reaction Conditions	8 μ M; 72h
Applications	18 α -Glycyrrhetic acid inhibited cell growth.
Animal experiment [2]:	
Animal models	BALB/c mice
Preparation Method	18 α -Glycyrrhetic acid solution was prepared in DMSO. A concentration of 10mg 18 α -Glycyrrhetic acid per kilogram of each mouse was used to be intraperitoneally injected in test group every other day within 30 consecutive days. The mice in the control group were treated with 2% DMSO (equal to the test group) as a vehicle in PBS. The expression of co-stimulatory molecules CD40 and MHC-II on isolated dendritic cells from mice were investigated by flow cytometry.
Dosage form	10mg/kg; 30d; q2d; i.p.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

Applications	18 α -Glycyrrhetic acid significantly decreased the expression of CD40 and MHC-II molecules.
References:[1] Zong L, Qu Y, Xu M Y, et al. 18 α -Glycyrrhetic acid extracted from <i>Glycyrrhiza radix</i> inhibits proliferation and promotes apoptosis of the hepatic stellate cell line [J]. Journal of digestive diseases, 2013, 14(6): 328-336.[2] Ebrahimnezhad S, Amirghofran Z, Karimi M H. Decline in immunological responses mediated by dendritic cells in mice treated with 18 α -glycyrrhetic acid [J]. Immunological investigations, 2016, 45(3): 191-204.	

Background

18 α -Glycyrrhetic acid is a bioactive triterpenoid mainly derived from *Glycyrrhiza radix*. 18 α -Glycyrrhetic acid has various pharmacological activities such as anti-oxidative and anti-inflammatory^[1]. 18 α -Glycyrrhetic acid can inhibit oxidative stress via activating Nrf-2 signaling^[1].

18 α -Glycyrrhetic acid (8 μ M; 72h) inhibited cell growth in Human HSC cells^[2]. 18 α -Glycyrrhetic acid (4 μ g/mL; 24h) significantly reduced ROS (reactive oxygen species) production in fructose-treated HK2 cells^[3]. 18 α -Glycyrrhetic acid (50 μ M; 48h) decreased the cell viability in Caco-2 cells^[4]. 18 α -Glycyrrhetic acid (2 μ g/mL; 24h) increased the cell viability in mitomycin C-induced HFL-1 cells^[1].

18 α -Glycyrrhetic acid (10mg/kg; 30d; q2d; i.p.) significantly decreased the expression of CD40 and MHC-II molecules in BALB/c mice^[5]. 18 α -Glycyrrhetic acid (12.5, 25 and 50mg/kg; 12 weeks; p.o.) alleviated fructose-induced renal injury in C57BL/6 mice^[3].

References:

- [1] Lefaki M, Papaevgeniou N, Tur J A, et al. The dietary triterpenoid 18 α -glycyrrhetic acid protects from MMC-induced genotoxicity through the ERK/Nrf2 pathway [J]. Redox biology, 2020, 28: 101317.
- [2] Zong L, Qu Y, Xu M Y, et al. 18 α -Glycyrrhetic acid extracted from *Glycyrrhiza radix* inhibits proliferation and promotes apoptosis of the hepatic stellate cell line [J]. Journal of digestive diseases, 2013, 14(6): 328-336.
- [3] Cheng X, Qiu L, Wang F. 18 α -Glycyrrhetic acid (GA) ameliorates fructose-induced nephropathy in mice by suppressing oxidative stress, dyslipidemia and inflammation [J]. Biomedicine &

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

pharmacotherapy, 2020, 125: 109702.

[4] Malekinejad M, Pashaei M R, Malekinejad H. 18 β -Glycyrrhetic acid altered the intestinal permeability in the human Caco-2 monolayer cell model [J]. European journal of nutrition, 2022, 61(7): 3437-3447.

[5] Ebrahimnezhad S, Amirghofran Z, Karimi M H. Decline in immunological responses mediated by dendritic cells in mice treated with 18 α -glycyrrhetic acid [J]. Immunological investigations, 2016, 45(3): 191-204.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA