

Product Data Sheet

Product Name: AG 825
Cat. No.: GC13168

Chemical Properties

Cas. No. 149092-50-2

Chemical Name (E)-3-(3-((benzo[d]thiazol-2-ylthio)methyl)-4-hydroxy-5-methoxyphenyl)-2-cyanoacrylamide

SMILES OC(C(CSC1=NC2=CC=CC=C2S1)=CC(/C=C(C#N)/C(N)=O)=C3)=C3OC


Formula $C_{19}H_{15}N_3O_3S_2$ M.Wt 397.47

Solubility DMF: 30 mg/ml, DMSO: 30 mg/ml, DMSO: PBS (pH 7.2)(1:3): 0.1 mg/ml, Ethanol: 0.2 mg/ml
Storage Store at -20°C

General tips For obtaining a higher solubility, please warm the tube at 37 °C and shake it in the ultrasonic bath for a while. Stock solution can be stored below -20°C for several months.

Shipping Condition Evaluation sample solution: ship with blue ice. All other available size: ship with RT, or blue ice upon request.

Structure



Background

AG 825 is a selective inhibitor of ErbB2 with IC₅₀ values of 0.15 and 19 μM for ErbB2 and ErbB1 respectively [1].

Receptor tyrosine-protein kinase erbB-2 is a member of the human epidermal growth factor receptor family and promotes cell proliferation.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

AG 825 is a selective ErbB2 inhibitor. AG825 inhibited autophosphorylation of HER2, HER1-2 and HER1 with IC₅₀ values of 0.15, 0.35 and 19 μ M, respectively [1]. In non-small cell lung cancer (NSCLC) cell lines expressing high-p185neu, the HER-2/neu gene product, AG825 significantly increased the chemosensitivities. However, AG825 with high concentrations inhibited the drug-induced G2 arrest and the activation of phosphorylated p34cdc2 [2]. In androgen-independent prostate cancer (PCa) cell lines C4 and C4-2, AG825 was toxic in a dose- and time-dependent way and inhibited phosphoactivation of HER-2/neu and its down-regulation. Also, AG825 induced an imbalance between p38 mitogen-activated protein kinase activation and extracellular signal-regulated kinase 1/2, which then led to p38-dependent apoptosis [3].

References:

- [1]. Osherov N, Gazit A, Gilon C, et al. Selective inhibition of the epidermal growth factor and HER2/neu receptors by tyrphostins. J Biol Chem, 1993, 268(15): 11134-11142.
- [2]. Tsai CM, Levitzki A, Wu LH, et al. Enhancement of chemosensitivity by tyrphostin AG825 in high-p185(neu) expressing non-small cell lung cancer cells. Cancer Res, 1996, 56(5): 1068-1074.
- [3]. Murillo H, Schmidt LJ, Tindall DJ. Tyrphostin AG825 triggers p38 mitogen-activated protein kinase-dependent apoptosis in androgen-independent prostate cancer cells C4 and C4-2. Cancer Res, 2001, 61(20): 7408-7412.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA