

Product Data Sheet

Product Name: Troglitazone

Cat. No.: GC15272

Chemical Properties

Cas. No. 97322-87-7

Chemical Name 5-[[4-[(6-hydroxy-2,5,7,8-tetramethyl-3,4-dihydrochromen-2-yl)methoxy]phenyl]methyl]-1,3-thiazolidine-2,4-dione

SMILES CC1=C(C(=C2CCC(OC2=C1C)(C)COC3=CC=C(C=C3)CC4C(=O)NC(=O)S4)C)O


Formula $C_{24}H_{27}NO_5S$ M.Wt 441.54

Solubility ≥ 20.9 mg/mL in DMSO, ≥ 3.34 mg/mL in EtOH with ultrasonic and warming
Storage Store at -20°C

General tips For obtaining a higher solubility, please warm the tube at 37°C and shake it in the ultrasonic bath for a while. Stock solution can be stored below -20°C for several months.

Shipping Condition Evaluation sample solution: ship with blue ice. All other available size: ship with RT, or blue ice upon request.

Structure



Protocol

Cell experiment [1]:

Cell lines C4-2 cells (human prostate cancer cell lines)

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

Cells were treated with Troglitazone 45μM for 72h, the percentage of cells within Preparation Method the different phases of the cell cycle was determined using flow cytometry, and the level of DNA fragmentation was measured using a Cell Death ELISA.

Reaction
Conditions 45μM; 72h

Applications Troglitazone treatment induced G₀/G₁ phase cell cycle arrest and increased the proportion of subG₀ phase cells and DNA fragmentation levels.

Animal experiment [2]:

Animal models Balb/c male mice bearing MIA Paca2 xenografts

Balb/c male mice were subcutaneously inoculated in the back with MIA Paca2 cells (5×10^6 cells) 14 days before the start of Troglitazone administration. Mice were then orally administered 200mg/kg of Troglitazone daily for 5 weeks.

Preparation Method Tumor volume was measured bi-dimensionally and volume was calculated by the formula $(\text{length} \times \text{width}^2) \times 0.5$, and mouse body weights were monitored throughout the experiment.

Dosage form 200mg/kg/day; 5 weeks; p.o.

Applications Troglitazone exhibited inhibitory effects on tumor growth in the MIA Paca2 xenograft model, the body weights of mice were not affected compared to those of the vehicle administration group.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

References:

- [1] AKINYEKE T O, STEWART L V. Troglitazone suppresses c-Myc levels in human prostate cancer cells via a PPAR γ -independent mechanism[J]. Cancer Biology & Therapy, 2011, 11(12): 1046-1058.
- [2] FUJITA M, HASEGAWA A, YAMAMORI M, et al. In vitro and in vivo cytotoxicity of troglitazone in pancreatic cancer[J]. Journal of Experimental & Clinical Cancer Research, 2017, 36(1): 91.

Background

Troglitazone is an orally active agonist of the peroxisome proliferator-activated receptor γ (PPAR γ), with EC₅₀ values of 550nM and 780nM for human and murine receptors, respectively [1]. PPAR γ is a widely expressed ligand-regulated transcription factor that controls the expression of genes involved in inflammation, redox balance, neurotrophic factor production, insulin sensitivity, and lipid and glucose metabolism[2]. By activating PPAR γ , Troglitazone improves insulin resistance and regulates glucose and lipid metabolism, making it commonly used for the treatment of type II diabetes[3].

In vitro, treatment of mitochondria isolated from rat liver with Troglitazone (10, 50 μ M) for 5min induced mitochondrial swelling[4]. Treatment of C4-2 cells with Troglitazone (45 μ M) for 72h induced G₀/G₁ phase cell cycle arrest and increased the proportion of cells in the subG₀ phase and the level of DNA fragmentation[5]. Treatment of human bladder cancer T24 and EJ cells with Troglitazone (25, 50 μ M) for 24h increased the ratio of light chain 3 (LC3)-II to actin in a time- and concentration-dependent manner[6].

In vivo, oral administration of Troglitazone (200mg/kg/day) for 5 weeks to male Balb/c mice bearing MIA Paca2 xenograft tumors significantly suppressed tumor growth without a significant effect on mouse body weight[7]. A single intraperitoneal injection of Troglitazone (300mg/kg) in female BALB/c mice significantly increased plasma alanine aminotransferase (ALT) and aspartate aminotransferase (AST) levels after 6h, and this treatment induced pericentral hepatocyte necrosis within 24h[8].

References:

- [1] WILSON T M, BROWN P J, STERNBACH D D, et al. The PPARs: from orphan receptors to drug discovery[J]. Journal of Medicinal Chemistry, 2000, 43(4): 527-550.
- [2] CAI W, YANG T, LIU H, et al. Peroxisome proliferator-activated receptor γ (PPAR γ): A master

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

- gatekeeper in CNS injury and repair[J]. Progress in Neurobiology, 2018, 163: 27-58.
- [3] PLOSKER G L, FAULDS D. Troglitazone: a review of its use in the management of type 2 diabetes mellitus[J]. Drugs, 1999, 57(3): 409-438.
- [4] OKUDA T, NORIOKA M, SHITARA Y, et al. Multiple mechanisms underlying troglitazone-induced mitochondrial permeability transition[J]. Toxicology and Applied Pharmacology, 2010, 248(3): 242-248.
- [5] AKINYEKE T O, STEWART L V. Troglitazone suppresses c-Myc levels in human prostate cancer cells via a PPAR γ -independent mechanism[J]. Cancer Biology & Therapy, 2011, 11(12): 1046-1058.
- [6] YAN S, YANG X, CHEN T, et al. The PPAR γ agonist Troglitazone induces autophagy, apoptosis and necroptosis in bladder cancer cells[J]. Cancer Gene Therapy, 2014, 21(5): 188-193.
- [7] FUJITA M, HASEGAWA A, YAMAMORI M, et al. In vitro and in vivo cytotoxicity of troglitazone in pancreatic cancer[J]. Journal of Experimental & Clinical Cancer Research, 2017, 36(1): 91.
- [8] JIA R, ODA S, TSUNEYAMA K, et al. Establishment of a mouse model of troglitazone-induced liver injury and analysis of its hepatotoxic mechanism[J]. Journal of Applied Toxicology, 2019, 39(11): 1541-1556.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA