

Product Data Sheet

Product Name: BGJ398 (Infigratinib)

Cat. No.: GC10055

Chemical Properties

Cas. No. 872511-34-7

Chemical Name 3-(2,6-dichloro-3,5-dimethoxyphenyl)-1-[6-[4-(4-ethylpiperazin-1-yl)anilino]pyrimidin-4-yl]-1-methylurea

SMILES CCN1CCN(CC1)C2=CC=C(C=C2)NC3=CC(=NC=N3)N(C)C(=O)NC4=C(C(=CC(=C4Cl)OC)OC)Cl


Formula $C_{26}H_{31}Cl_2N_7O_3$ M.Wt 560.48

Solubility $\geq 7\text{mg/mL}$ in DMSO with gentle warming Storage Store at -20°C

General tips For obtaining a higher solubility, please warm the tube at 37°C and shake it in the ultrasonic bath for a while. Stock solution can be stored below -20°C for several months.

Shipping Condition Evaluation sample solution: ship with blue ice. All other available size: ship with RT, or blue ice upon request.

Structure



Protocol

Cell experiment [1]:

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

Cell lines SKOV3ip1 cells

Preparation Method To measure cell viability of spheroid cultures, SKOV3ip1 cells (5×10^4) were seeded in ultra-low attachment 24-well plates and incubated overnight. BGJ398 were then added, and the cells were incubated for an additional 72 hours. Cells were harvested from each well and transferred to a regular 24-well plate for an additional 12 hours. Attached, live cells were stained with 0.2% crystal violet solution for 20 minutes with gentle agitation. The stained cells were washed with distilled water. For colorimetric analysis, the crystal violet dye was extracted with 1% SDS/PBS, and absorbance was measured at 570nm using an EMax PLUS microplate reader.

Reaction Conditions 0.3125-5 μ M; 72h

Applications The survival rate of SKOV3ip1 cells in spheroid culture was significantly reduced after BGJ398 treatment.

Animal experiment [2]:

Animal models TMK-1 cells xenograft mice model

Preparation Method Cancer cells (1×10^6) were injected subcutaneously into the right flank of 6- to 8-week-old athymic nude mice. Mice were randomly assigned to either a control or treatment group. When tumors reached approximately 80mm³, BGJ398 (10mg/kg, administered by oral gavage) was initiated. Tumor diameters were measured, and volumes were calculated ($\text{width}^2 \times \text{length} \times 0.5$). Tumors were excised, weighed, and processed for subsequent experiments.

Dosage form 10mg/kg; po; 20d

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

Applications Treatment with BGJ398 significantly inhibited tumor growth.

References:

- [1]. Cha H J, Choi J H, Park I C, et al. Selective FGFR inhibitor BGJ398 inhibits phosphorylation of AKT and STAT3 and induces cytotoxicity in sphere-cultured ovarian cancer cells[J]. International Journal of Oncology, 2017, 50(4): 1279-1288.
- [2]. Schmidt K, Moser C, Hellerbrand C, et al. Targeting fibroblast growth factor receptor (FGFR) with BGJ398 in a gastric cancer model[J]. Anticancer Research, 2015, 35(12): 6655-6665.

Background

BGJ398 (Infigratinib) is an oral, ATP-competitive pan-FGFR tyrosine kinase inhibitor (FGFR1: IC₅₀ = 0.9nM; FGFR2: IC₅₀ = 1.4nM; FGFR3: IC₅₀ = 1nM; FGFR4: IC₅₀ = 60nM) [1]. BGJ398 targets the fibroblast growth factor receptors FGFR1, FGFR2, FGFR3, and FGFR4, thereby blocking signal transduction mediated by these receptors and inhibiting FGFR-dependent tumor cell proliferation and angiogenesis [2-3]. BGJ398 is primarily used to treat dwarfism in children [4].

In SKOV3ip1 cells, the survival rate of cell in spheroid culture was significantly reduced after BGJ398 (0.3125-5μM; 72h) treatment [5]. In UMUC-14 cells, the combination of paclitaxel and BGJ398 (0-10μM; 72h) can produce a synergistic effect in promoting cell apoptosis [6].

In TMK-1 cells xenograft mice model, treatment with BGJ398 (10mg/kg; po; 20d) significantly inhibited tumor growth [7]. In Snu16R-derived xenograft mice model, knockdown of BAG3 enhances the anti-tumor effect induced by BGJ398 (30mg/kg; po; 4 weeks) [8].

References:

- [1]. Guagnano V, Furet P, Spanka C, et al. Discovery of 3-(2, 6-dichloro-3, 5-dimethoxy-phenyl)-1-{6-[4-(4-ethyl-piperazin-1-yl)-phenylamino]-pyrimidin-4-yl}-1-methyl-urea (NVP-BGJ398), a potent and selective inhibitor of the fibroblast growth factor receptor family of receptor tyrosine kinase[J]. Journal of medicinal chemistry, 2011, 54(20): 7066-7083.
- [2]. Szymczyk J, Sluzalska K D, Materla I, et al. FGF/FGFR-dependent molecular mechanisms underlying anti-cancer drug resistance[J]. Cancers, 2021, 13(22): 5796.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

- [3]. Agrawal S, Maity S, AlRaawi Z, et al. Targeting drugs against fibroblast growth factor (s)-induced cell signaling[J]. Current drug targets, 2021, 22(2): 214-240.
- [4]. Höglér W, Ward L M. New developments in the management of achondroplasia[J]. Wiener Medizinische Wochenschrift, 2020, 170(5): 104-111.
- [5]. Cha H J, Choi J H, Park I C, et al. Selective FGFR inhibitor BGJ398 inhibits phosphorylation of AKT and STAT3 and induces cytotoxicity in sphere-cultured ovarian cancer cells[J]. International Journal of Oncology, 2017, 50(4): 1279-1288.
- [6]. Kim S H, Ryu H, Ock C Y, et al. BGJ398, a pan-FGFR inhibitor, overcomes paclitaxel resistance in urothelial carcinoma with FGFR1 overexpression[J]. International journal of molecular sciences, 2018, 19(10): 3164.
- [7]. Schmidt K, Moser C, Hellerbrand C, et al. Targeting fibroblast growth factor receptor (FGFR) with BGJ398 in a gastric cancer model[J]. Anticancer Research, 2015, 35(12): 6655-6665.
- [8]. Li K, Deng X, Feng G, et al. Knockdown of Bcl-2-associated athanogene-3 can enhance the efficacy of BGJ398 via suppressing migration and inducing apoptosis in gastric cancer[J]. Digestive Diseases and Sciences, 2021, 66(9): 3036-3044.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA