

Product Data Sheet

Product Name: Fulvestrant (ICI 182,780)

Cat. No.: GC18000

Chemical Properties

Cas. No. 129453-61-8

Chemical Name (7R,8R,9S,13S,14S,17S)-13-methyl-7-[9-(4,4,5,5,5-pentafluoropentylsulfinyl)nonyl]-6,7,8,9,11,12,14,15,16,17-decahydrocyclopenta[a]phenanthrene-3,17-diol

SMILES CC12CCC3C(C1CCC2O)C(CC4=C3C=CC(=C4)O)CCCCCCCCCS(=O)CCCC(C(F)(F)F)(F)F


Formula $C_{32}H_{47}F_5O_3S$ M.Wt 606.77

Solubility $\geq 30.35\text{mg/mL}$ in DMSO Storage Store at -20°C

General tips For obtaining a higher solubility, please warm the tube at 37°C and shake it in the ultrasonic bath for a while. Stock solution can be stored below -20°C for several months.

Shipping Condition Evaluation sample solution: ship with blue ice All other available size: ship with RT, or blue ice upon request.

Structure



Protocol

Cell experiment [1]:

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

Cell lines	Bats-72 cell line (estrogen receptor-negative MDR cell line)
Preparation method	Cells were exposed to designated concentrations of doxorubicin with or without co-treatment of fulvestrant for 72 h, the cell viabilities were determined.
Reaction Conditions	1-10 μ M;72 h
Applications	Fulvestrant sensitizes doxorubicin-induced cytotoxicity in estrogen receptor-negative MDR cell line Bats-72.
Animal experiment [2]:	
Animal models	Balb/C-nu/nu mice (MCF7 cells model)
Preparation method	Mice were treated with fulvestrant in addition to the estradiol exposure.
Dosage form	5 mg/mouse; s.c.; twice per week
Applications	Fulvestrant significantly inhibited macrophage and neutrophil infiltration in mice cancer model.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

References:

- [1]. Huang Y, Jiang D, et, al. Fulvestrant reverses doxorubicin resistance in multidrug-resistant breast cell lines independent of estrogen receptor expression. *Oncol Rep.* 2017 Feb;37(2):705-712. doi: 10.3892/or.2016.5315. Epub 2016 Dec 14. PMID: 28000875; PMCID: PMC5355712.
- [2]. Abrahamsson A, Rodriguez GV, et, al. Fulvestrant-Mediated Attenuation of the Innate Immune Response Decreases ER+ Breast Cancer Growth In Vivo More Effectively than Tamoxifen. *Cancer Res.* 2020 Oct 15;80(20):4487-4499. doi: 10.1158/0008-5472.CAN-20-1705. Epub 2020 Aug 27. PMID: 32855207.

Background

Fulvestrant is a selective estrogen receptor (ER) antagonist. It binds, blocks and degrades estrogen receptor, then inhibits estrogen receptor(ER)-mediated transcriptional activity with an IC_{50} of 9.4 nM [1-3].

Fulvestrant(3 μ M ;1 month) increases the sensitivity of H1975 NSCLC cells to gefitinib^[4]. Fulvestrant(1-10 μ M;72 h) strongly sensitized doxorubicin-induced cytotoxicity in MDR cell lines^[5].

Fulvestrant(5 mg/mouse; s.c.; twice per week) significantly inhibited macrophage and neutrophil infiltration in mice cancer model. Fulvestrant decreases ER⁺ breast cancer growth in the presence of physiologic levels of estradiol in human breast cancer in nude mice^[6]. Fulvestrant (5mg/mouse; twice weekly; s.c.) in combination with tamoxifen enhanced tumor regression^[7].

References:

- [1]. Vergote I, Robertson JF. Fulvestrant is an effective and well-tolerated endocrine therapy for postmenopausal women with advanced breast cancer: results from clinical trials. *Br J Cancer.* 2004 Mar;90 Suppl 1(Suppl 1):S11-4. doi: 10.1038/sj.bjc.6601631. PMID: 15094759; PMCID: PMC2750769.
- [2]. Osborne CK, Wakeling A, et, al. Fulvestrant: an oestrogen receptor antagonist with a novel

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

mechanism of action. Br J Cancer. 2004 Mar;90 Suppl 1(Suppl 1):S2-6. doi: 10.1038/sj.bjc.6601629. PMID: 15094757; PMCID: PMC2750773.

[3]. Dowsett M, Nicholson RI, et.al. Biological characteristics of the pure antiestrogen fulvestrant: overcoming endocrine resistance. Breast Cancer Res Treat. 2005;93 Suppl 1:S11-8. doi: 10.1007/s10549-005-9037-3. PMID: 16247595.

[4]. Shen H, Liu J, et.al. Fulvestrant increases gefitinib sensitivity in non-small cell lung cancer cells by upregulating let-7c expression. Biomed Pharmacother. 2014 Apr;68(3):307-13. doi: 10.1016/j.biopha.2013.10.007. Epub 2013 Nov 7. PMID: 24268810.

[5]. Huang Y, Jiang D, et.al. Fulvestrant reverses doxorubicin resistance in multidrug-resistant breast cell lines independent of estrogen receptor expression. Oncol Rep. 2017 Feb;37(2):705-712. doi: 10.3892/or.2016.5315. Epub 2016 Dec 14. PMID: 28000875; PMCID: PMC5355712.

[6]. Abrahamsson A, Rodriguez GV, et.al. Fulvestrant-Mediated Attenuation of the Innate Immune Response Decreases ER+ Breast Cancer Growth In Vivo More Effectively than Tamoxifen. Cancer Res. 2020 Oct 15;80(20):4487-4499. doi: 10.1158/0008-5472.CAN-20-1705. Epub 2020 Aug 27. PMID: 32855207.

[7]. Mishra AK, Abrahamsson A, et.al. Fulvestrant inhibits growth of triple negative breast cancer and synergizes with tamoxifen in ER α positive breast cancer by up-regulation of ER β . Oncotarget. 2016 Aug 30;7(35):56876-56888. doi: 10.18632/oncotarget.10871. PMID: 27486755; PMCID: PMC5302959.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA