

Product Data Sheet

Product Name: Neobavaisoflavone

Cat. No.: GC30085

Chemical Properties

Cas. No. 41060-15-5

SMILES O=C1C(C2=CC=C(O)C(C/C=C(C)\C)=C2)=COC3=CC(O)=CC=C13

Formula $C_{20}H_{18}O_4$ M.Wt 322.35

Solubility DMSO : ≥ 31 mg/mL (96.17 mM) Storage Store at -20°C

General tips For obtaining a higher solubility , please warm the tube at 37°C and shake it in the ultrasonic bath for a while. Stock solution can be stored below -20°C for several months.

Shipping Condition Evaluation sample solution : ship with blue ice All other available size: ship with RT , or blue ice upon request.


Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

Structure



Background

Neobavaisoflavone, an isoflavone isolated from *Psoralea corylifolia*, has striking anti-inflammatory and anti-cancer effects. IC₅₀ value: 42.93 μ M (toward CCRF-CEM cells); 114.64 μ M [against HCT116 (p53(+/+)) cells] [2] Target: In vitro: In the cancer cells, neobavaisoflavone sensitizes human U373MG glioma cells to TRAIL-mediated apoptosis; upregulated DR5 expression; induced TRAIL-mediated apoptosis in human glioma cells by suppressing migration and invasion, and by inhibiting anoikis resistance [1]. In cancer cell lines, neobavaisoflavone is selectively active, and IC₅₀ values below 115 μ M were obtained on 6/9 cell lines, with values ranging from 42.93 μ M (toward CCRF-CEM cells) to 114.64 μ M [against HCT116 (p53(+/+)) cells] [2]. In vivo:

[1]. Kim YJ, et al. Neobavaisoflavone sensitizes apoptosis via the inhibition of metastasis in TRAIL-resistant human glioma U373MG cells. *Life Sci.* 2014 Jan 30;95(2):101-7. [2]. Kuete V, Activity of three cytotoxic isoflavonoids from *Erythrina excelsa* and *Erythrina senegalensis* (neobavaisoflavone, sigmoidin H and isoneorautenol) toward multi-factorial drug resistant cancer cells. *Phytomedicine*.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

2014 Apr 15;21(5):682-8. [3]. Ming-Jaw Don, et al. Neobavaisoflavone stimulates osteogenesis via p38-mediated up-regulation of transcription factors and osteoid genes expression in MC3T3-E1 cells. Phytomedicine Volume 19, Issue 6, 15 April 2012, Pages 551-561

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA