

Product Data Sheet

Product Name: PFK-015

Cat. No.: GC16993

Chemical Properties

Cas. No. 4382-63-2

SMILES O=C(C1=CC=NC=C1)/C([H])=C([H])/C2=NC3=CC=CC=C3C=C2


Formula $C_{17}H_{12}N_2O$ M.Wt 260.29

Solubility $\geq 13\text{mg/mL}$ in DMSO Storage Store at -20°C

General tips For obtaining a higher solubility, please warm the tube at 37°C and shake it in the ultrasonic bath for a while. Stock solution can be stored below -20°C for several months.

Shipping Condition Evaluation sample solution: ship with blue ice. All other available size: ship with RT, or blue ice upon request.

Structure



Protocol

Cell experiment [1]:

Cell lines Peripheral blood mononuclear cells

Preparation Method Peripheral blood mononuclear cells supplemented with 40ng/ml macrophage colony-stimulating factor (M-CSF) was used as a differentiation medium. To examine the role of PFKFB3 in macrophage polarization, PFKFB3 inhibitor PFK-015 ($10\mu\text{M}$) was used.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

Reaction Conditions 10 μ M; 80min

Applications Treatment with PFK-015 during IFN- γ -induced macrophage polarization reduced the expression of M1 surface markers CD80, iNOS, and HLA-DR.

Animal experiment [2]:

Animal models Xenograft mode

Preparation Method LL cells were obtained from DMEM medium containing 10% FCS in the exponential growth phase. After washing twice, the cells were resuspended in PBS buffer. 0.1mL of cell suspension (1×10^6 cells) was subcutaneously injected into C57Bl6 mice. Mice carrying xenograft tumors (150–200mg) were randomly assigned to four groups: DMSO group, PFK-015 group (25mg/kg, intraperitoneal injection, every 3 days for 4 doses), irinotecan group (70mg/kg, intravenous injection, every 3 days for 4 doses), temozolomide group (30mg/kg, oral administration, 3 consecutive days followed by a 3-day break), and gemcitabine group (100mg/kg, every 3 days for 4 doses). Tumor size was measured daily using microcalipers.

Dosage form 25mg/kg; iv; every 3 days for 4 doses

Applications PFK-015 inhibits the growth of Lewis lung carcinoma (LLC) xenografts by mediating the depletion of tumor-associated fructose-2,6-bisphosphate (F26BP), which consequently triggers apoptotic signaling and suppresses pulmonary metastasis of the subcutaneous tumors.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

References:

- [1]. Chen R, Wang J, Dai X, et al. Augmented PFKFB3-mediated glycolysis by interferon- γ promotes inflammatory M1 polarization through the JAK2/STAT1 pathway in local vascular inflammation in Takayasu arteritis[J]. Arthritis research & therapy, 2022, 24(1): 266.
- [2]. Clem B F, O'Neal J, Tapolsky G, et al. Targeting 6-phosphofructo-2-kinase (PFKFB3) as a therapeutic strategy against cancer[J]. Molecular cancer therapeutics, 2013, 12(8): 1461-1470.

Background

PFK-015 is a potent and selective inhibitor of PFKFB3 with an IC₅₀ value of 110nM^[1]. As a key regulator of glycolytic metabolism, PFK-015 disrupts fructose-2,6-bisphosphate production and demonstrates significant anti-proliferative effects across multiple tumor cell lines^[2]. The compound also induces autophagy and exhibits synergistic activity with chemotherapeutic agents in vivo^[3].

In Peripheral blood mononuclear cells treatment with PFK-015(10 μ M; 80min) during IFN- γ -induced macrophage polarization reduced the expression of M1 surface markers CD80, iNOS, and HLA-DR^[4]. In the 786-O and OS-RC-2 cell lines, the expression levels of glycolytic regulatory enzymes, key enzymes, and PD-L1 decreased after the addition of the PFKFB3 inhibitor PFK-015(10 μ M; 24h)^[5].

In Lewis lung carcinoma (LLC)-bearing mice, PFK-015(25mg/kg; iv; every 3 days for 4 doses) inhibits the growth of Lewis lung carcinoma (LLC) xenografts by mediating the depletion of tumor-associated fructose-2,6-bisphosphate (F26BP), which consequently triggers apoptotic signaling and suppresses pulmonary metastasis of the subcutaneous tumors^[6]. In the Hu-NOG mouse model, PFK-015(25mg/kg; ip; 21d) combined with PD-1 mAb therapy significantly inhibits tumor growth in human PBMC-engrafted NOG mice^[7].

References:

- [1].Clem B, Tapolsky G H, O'Neal J, et al. Characterization of a novel small molecule antagonist of 6-phosphofructo-2-kinase that suppresses glucose metabolism and tumor growth[J]. Cancer Research, 2011, 71(8_Supplement): 2825-2825.
- [2].Ling X, Liu L, Jiang A, et al. PFKFB3 promotes endometriosis cell proliferation via enhancing the

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA

Product Data Sheet

- protein stability of β -catenin[J]. Molecular and Cellular Endocrinology, 2024, 579: 112083.
- [3].Brooke D G, van Dam E M, Watts C K W, et al. Targeting the Warburg Effect in cancer; relationships for 2-arylpyridazinones as inhibitors of the key glycolytic enzyme 6-phosphofructo-2-kinase/2, 6-bisphosphatase 3 (PFKFB3)[J]. Bioorganic & medicinal chemistry, 2014, 22(3): 1029-1039.
- [4].Chen R, Wang J, Dai X, et al. Augmented PFKFB3-mediated glycolysis by interferon- γ promotes inflammatory M1 polarization through the JAK2/STAT1 pathway in local vascular inflammation in Takayasu arteritis[J]. Arthritis research & therapy, 2022, 24(1): 266.
- [5].Yu Y, Liang Y, Li D, et al. Glucose metabolism involved in PD-L1-mediated immune escape in the malignant kidney tumour microenvironment[J]. Cell Death Discovery, 2021, 7(1): 15.
- [6].Clem B F, O'Neal J, Tapolsky G, et al. Targeting 6-phosphofructo-2-kinase (PFKFB3) as a therapeutic strategy against cancer[J]. Molecular cancer therapeutics, 2013, 12(8): 1461-1470.
- [7].Zheng J B, Wong C W, Liu J, et al. Glucose metabolism inhibitor PFK-015 combined with immune checkpoint inhibitor is an effective treatment regimen in cancer[J]. Oncoimmunology, 2022, 11(1): 2079182.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: (909) 407-4943 Fax: (626) 353-8530 E-mail: tech@glpbio.com

Address: 10292 Central Ave. #205, Montclair, CA, USA